

Compliance to Treatment and Quality of Life of Sudanese Patients with Heart Failure

Mugahed Ali Ahmed AL-khadher^{1, *}, Imad Fadl-Elmula²,
Waled Amen Mohammed Ahmed³

¹Medical-Surgical Nursing Department, Sudan International University, PhD Nursing Student of Medical-Surgical Nursing, Faculty of Nursing Al Neelain University, Khartoum, Sudan

²Department of Pathology, Nahda College, Khartoum, Sudan

³Faculty of Applied Medical Sciences, Albaha University, Al-Baha, Kingdom of Saudi Arabia

Abstract

Background: Heart failure is known to decrease the quality of life, especially in non-compliance patients with regards to medications and life style changes. *Objective:* The present study aimed to determine the level of compliance to treatment and quality of life of Sudanese patients with heart failure. *Methods:* This descriptive study was conducted on 76 patients with heart failure admitted to the Sudan Heart Institute. Demographic and clinical data including the compliance (medication, sodium restriction, fluid restriction, daily weights, exercises, and appointment-keeping) were collected. The quality of life was measured using the Minnesota living with heart failure Questionnaire. The data were collected from all patients and the analyzed using SPSS version 22 software. *Results:* Heart failure patients showed low compliance ranged between 11.84% and 75% of which the highest compliance was to medication (75%) followed by the follow-up appointments (71.05%), and the lowest compliances were to the fluids restrictions (11.84%), the weight monitoring (17.10%), regular exercise (21.05%), and the sodium restriction (27.6%). Quality of life score ranged between 62-97 score and the Mean (SD) 83.6 (7.82) which revealed of poor quality of life in most of Sudanese patients with heart failure involved in the present study. *Conclusion:* The study showed that patients with heart failure in Sudan have low compliance to treatment and poor quality of life.

Keywords

Heart Failure, Treatment Compliance, Quality of Life, Sudan

Received: April 23, 2015 / Accepted: May 3, 2015 / Published online: May 27, 2015

© 2015 The Authors. Published by American Institute of Science. This Open Access article is under the CC BY-NC license.

<http://creativecommons.org/licenses/by-nc/4.0/>

1. Introduction

Heart failure incidence increases with age, increase from approximately 20 per 1000 individuals with age 65 to 69-year-old to more than 80 per 1000 individuals aging 85-year-old. In fact few epidemiological data on heart failure in Sudan exists and the recognition of the disease as a major health issue remains questionable, the prevalent of heart failure accounts for 2.5% of the population, and hence it is one of the major causes of hospital mortality.

The WHO defined adherence as extent a person's behavior – taking drugs, following a diet, and/or executing lifestyle modifications, follow the agreed recommendations from a health care providers. Poor compliance “noncompliance” usually refers to patients' failure to follow health interventions as recommended by the health care provider, but it can also refer to the providers' failure to act according to practice guidelines or standards of care. The factors affecting the compliance could be divided into patient-related

* Corresponding author

E-mail address: malkhder2004@yahoo.com (M. A. A. AL-khadher)

factors, regimen-related factors, and health care providers-related factors.

Non-compliance to medications and diet contributes in many cases to worsening heart failure symptoms. The compliance to prescribe medications or other caregivers recommendations such as lifestyle changes is a widely acknowledged problem leading to hospitalization ((6-8). The non-compliance of HF patients is a major problem and remains to be a continuous source of concern for patients. It is mainly for diet and fluid, daily weight and exercises.

Quality of life (QOL) is defined as the individual's unique cognition and a way to express feelings about his/her health status. Moreover, QOL is a good predictor of mortality and the need for hospitalization (11-13). Patients in class II and III heart failure of New York Heart Association (NYHA) classification cannot normally do their daily activities.

Although, several studies on compliance of HF patients and their quality of life have been performed worldwide, to our knowledge this is the first ever study conducted in Sudanese HF patients, aimed to assess the compliance to treatment and quality of life in Sudanese patients with heart failure.

2. Materials and Methods

This descriptive study was conducted on patients with heart failure admitted to the Sudan Heart Institute. A total of 76 Sudanese HF patients were randomly selected from Sudan Heart Institute in Khartoum, January-March 2014. The patients participated were 20 years old or older, diagnosed as heart failure by the cardiologist at least a month, already started HF treatment, in class II or III heart failure of NYHA, and with ability to communicate.

The questionnaire consists of 36 questions of which 10 for demographic and clinical data, 5 questions for compliance, and 21 questions for quality of life. Demographic and clinical data were collected from medical records and/or by interviews. The demographic data included age, gender, educational level, and marital status, whereas clinical variables include left ventricular ejection fraction (EF), previous hospitalization in the past three months, and duration of HF.

Revised HF Compliance Questionnaire was used, on a five-point scale (1='never'; 2= seldom; 3= half of the time; 4 =mostly; 5='always'). the participant's compliance to medications, diet, fluid restriction, exercise, weight, and appointment keeping was evaluated by asking patients to rate their compliance of the last week (drugs, diet modifications, fluid restriction, and exercises), the last month (daily weighing), and the last 3 months (appointment keeping) before hospitalization. The patients were divided into two groups; either compliant or noncompliant (16-19). Patients

were considered 'overall compliant' the compliance with four or more of the six recommendations.

The quality of life data were collected and measured using the Minnesota Living with Heart Failure Questionnaire after translated to Arabic language. This instrument used most widely to evaluate quality of life in research studies (21-24). Which Contains 21 questions and overall score of 105 (5x21) with possible answers ranging from 0 (no) to 5 (very much), (0= no; 1= Very Little; 2= little; 3= moderate; 4= much; 5= very much). The final score is the sum points obtained for the 21 questions; it can therefore vary between 0 and 105. It evaluates how heart failure affects patients 'physical (8 questions), emotional (5 questions), and socioeconomic (8 questions) dimensions. The sum of responses reflects the overall effects of heart failure and treatments on individual's quality of life.

Data was presented using descriptive statistics including frequency, percentage, mean with standard deviation (SD) and P-value of ≤ 0.05 was considered statistically significant for relationship investigations. Ethical approval was obtained from Al Neelain Ethical committee at Al Neelain University.

3. Results

The study showed that out the 76 patients, 63.2% were male and 36.8% were female; the mean age was 61.4 ± 13.5 years. The education levels were 34.2% of patients were illiterate, 32.9% had completed primary school, 19.7% secondary school, and 13.2% had university graduation (Table 1).

Although the vast majority of the patients were chronic patients with diagnosis for more than 5 years, the participant ask to define what is the heart failure? Only 24% had basic conscious about their disease, the remaining 76% of patients had no idea what the heart failure is. Overall compliance among the patients was 28.95%, whereas 71.5% of the patients were classified as non-compliant. Of those compliance with medication was 75% and 70% compliance with appointment-keeping. In general most patients showed low compliance with diet restriction (27%), exercise (21%), weighing (17%), and fluid restriction (11%) (Table2).

The quality of life data showed that poor quality of life, the score ranged from 62-97 score /105, and the Mean (SD) quality of life was 3.2 (1.3) which revealed poor quality of life in most of Sudanese patients with heart failure involved in the present study. There is statistically significant in compliance and quality of life (p value= 0.002) in comparison with patients who is noncompliant. Also statistically significant with improved NYHA classification, LVEF and quality of life (p<0.001), others demographic and clinical data showed statistically insignificant (Table 3).

Table 1. Demographic and clinical variables of the study population (n=76) in Sudan.

Characteristic		Frequency	%
Age (years SD)	Mean (SD)		61.4 ±13.5
Sex	Male	48	63.2%
	Female	28	36.8 %
Marital status	Married	55	72.4%
	Single	8	10.5%
	Widowed	11	14.5%
	Divorced	2	2.6%
Employment Status	Employed	27	35%
	Unemployed	38	50.5%
	Retired	11	14.5 %
Educational Level	Non	26	34.%
	Primary	25	232.9%
	Secondary	15	19.7%
	University/college	10	13.2%
Duration Of Disease	Less than one year ago	38	50.0%
	One to three year ago	30	39.47%
	Four years and above ago	8	10.5 %
NYHA class	Class II	38	55.0 %
	Class III	33	43.4%
	Class IV	5	6.6 %
Ejection Fraction	mean (SD)	37 ±14	
Previous HF Emergency admission	No admission	21	27.6%
	One admission	29	38.2 %
	>1	26	34.2 %

Table 2. Compliance (Medications, diet, Fluid restriction, Exercise, weight, and appointments keeping) in Sudan.

How often	Compliant	Non-compliant
1. Do you take your medications exactly as directed?	(75%) 57	25%) 19
2. Do you weigh yourself daily? Or at least three times/week?	(17.10 %) 13	(82.89%) 63
3. Do you follow a low sodium diet?	(27.63%) 21	(72.36 %) 55
4. Do you avoid drinking excess fluids?	(11.84%) 9	(88.15 %) 67
5. Do you get regular exercise?	(21.05)% 16	(78.9%) 60
6. Do you Keep follow-up appointments?	(71.05) % 54	(28..9%) 22

Table 3. Quality of life of heart failure patients in Sudan (N=76).

Quality of life items	Mean	Std. Deviation
Causing swelling in your ankles or legs?	3.8026	1.11976
Making you sit or lie down to rest during the day?	3.5395	1.47369
Making your working around the house or yard difficult?	3.5132	1.21648
Making your going places away from home difficult?	3.8421	1.49713
Making your sleeping well at night difficult?	3.5395	1.30067
Making your relating to or doing things with your friends or family difficult?	3.8421	1.37649
Making your working to earn a living difficult?	3.6974	1.39542
Making your recreational pastimes, sports or hobbies difficult?	4.2368	1.00490
Making your sexual activities difficult?	4.5658	.86926
Making you eat less of the foods you like?	3.9737	1.49643
Making you tired, fatigued, or low on energy?	4.5263	1.02598
Making you stay in a hospital?	3.9079	1.23480
Costing you money for medical care?	4.1316	1.19267
Giving you side effects from treatments?	4.2895	.97729
Making you feel a loss of self-control in your life?	4.0395	1.18255
Making you worry?	4.2895	1.16408
Making your walking about or climbing stairs difficult?	4.0658	1.35976
Making you tired, fatigued, or low on energy?	4.1447	1.16280
Making you feel you are a burden to your family or friends?	3.9737	1.35621
Making it difficult for you to concentrate or remember things?	3.8553	1.50292
Making you feel depressed?	3.8553	1.19671
Mean ±SD	3.2±1.3	

4. Discussion

The patients' compliance in this study ranged between 11.84% and 75% of the patients. Although the differences in measurement instruments and differences in interventions, the result of the this study showed low compliance compared with other previous studies including knowledge of the patients about their illness, the hazard of high salt consumption, and the daily weighing.

Study done by Baghianimoghadam MH, et al, reported that the disease knowledge in Iranian patients reached 38%, whereas our result showed that 76% of HF Sudanese Patients lack essential knowledge of their disease or what the heart failure is. According to definition of 'overall compliance, the overall patients' compliance of the present study was 28% compared with the study conducted by van der wal in which the overall compliance reached 72% of patients with HF. In the same study compliance with medication (98.6%), appointment keeping, salt restriction (79%), fluid restriction (73%), exercise (39%), and weighing (35%) where all higher compared with the results of the present study. Also the compliance level of present study is lower than Evangelista study which found higher levels of compliance more than 90% for (follow-up appointments, medications, smoking, and alcohol cessation), low compliance dietary 71% and exercise recommendations 53%. Medication compliance in the present study result is similar to the study done by Kamlovi Yayhd which found %74.7 that compliance to medication.

The Minnesota living with heart failure questionnaire (MLWHFQ) showed that poor quality of life, the score ranged between 62-97 score /105, and the Mean (SD) quality of life was 83.6 (7.82) which revealed poor quality of life in most of Sudanese patients with heart failure involved in the present study.

It was also found that no correlation between age and quality of life (p value =0.925), this similar to study done by Kato N, et al, some studies found association between age and quality of life. We did not observe sex differences in quality of life (p value =0.99), which similar to study done by Heo S, et al 2007. But other studies have reported quality of life worse in female (30;31).

Also we found marital status had no influence on QOL in our subjects (p value =0.34), it is lower to study done by Luttk ML, which found differences in QoL between married patients and those living alone were most pronounced with regard to future expectations of QoL (6.5 vs 5.0, P=.00).

Our study shows there is statistically significant in duration of disease with QOL (p value =0.004), Also statistically significant with improved NYHA classification, LVEF and

Quality of life (p<0.001). This might be explained, partly, by the sedentary life style because of HF mostly effect elder people, lack of awareness of the importance of physical exercise, a culture that discourages physical exercise especially for females, the absence of safe public places where one could go walking and the hot of the weather in Sudan.

The strengths of this study include the study conducted in developing countries with limited resources and in short period of time. The study limitations were; the study was conducted as a descriptive study; interventional studies will yield more useful results if conducted on more sample with complete randomization all over the country.

In this study, the researcher found that total compliance was poor for HF Sudanese patients, compliance for drugs and appointments keeping were high but still in an unacceptable level. Compliance with diet, fluid restriction, activity and daily weighing was low. Also the study revealed that non-compliance negatively affects the quality of life of Sudanese HF patients. Based on result of present study, education and counselling are extremely needed to increased patients-knowledge about their disease, leading to more compliance and improvement of their quality of life.

References

- [1] Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE, Jr., Drazner MH, et al. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2013 Oct 15; 62:e147-e239.
- [2] Suliman A. The state of heart disease in Sudan. *Cardiovasc J Afr* 2011 Jul; 22:191-6.
- [3] De GS, Sabate E. Adherence to long-term therapies: evidence for action. *Eur J Cardiovasc Nurs* 2003 Dec;2 :323.
- [4] Kane S, Huo D, Aikens J, Hanauer S. Medication nonadherence and the outcomes of patients with quiescent ulcerative colitis. *Am J Med* 2003 Jan; 114:39-43.
- [5] Van der Wal MH, Jaarsma T, Moser DK, Veeger NJ, van Gilst WH, van Veldhuisen DJ. Compliance in heart failure patients: the importance of knowledge and beliefs. *Eur Heart J* 2006 Feb; 27:434-40.
- [6] Ghali JK, Kadakia S, Cooper R, Ferlinz J. Precipitating factors leading to decompensation of heart failure. Traits among urban blacks. *Arch Intern Med* 1988 Sep; 148:2013-6.
- [7] Michalsen A, Konig G, Thimme W. Preventable causative factors leading to hospital admission with decompensated heart failure. *Heart* 1998 Nov; 80:437-41.
- [8] Tsuyuki RT, McKelvie RS, Arnold JM, Avezum A, Jr., Barretto AC, Carvalho AC, et al. Acute precipitants of congestive heart failure exacerbations. *Arch Intern Med* 2001 Oct 22; 161:2337-42.

- [9] Lakdzaji S, Hassankhni H, Mohajjel AA, Khajegodary M, Salehi R. Effect of educational program on quality of life of patients with heart failure: a randomized clinical trial. *J Caring Sci* 2013 Mar; 2:11-8.
- [10] Gill TM, Feinstein AR. A critical appraisal of the quality of quality-of-life measurements. *JAMA* 1994 Aug 24; 272:619-26.
- [11] Hulsmann M, Berger R, Sturm B, Bojic A, Woloszczuk W, Bergler-Klein J, et al. Prediction of outcome by neurohumoral activation, the six-minute walk test and the Minnesota Living with Heart Failure Questionnaire in an outpatient cohort with congestive heart failure. *Eur Heart J* 2002 Jun; 23:886-91.
- [12] Konstam V, Salem D, Pouleur H, Kostis J, Gorkin L, Shumaker S, et al. Baseline quality of life as a predictor of mortality and hospitalization in 5,025 patients with congestive heart failure. SOLVD Investigations. Studies of Left Ventricular Dysfunction Investigators. *Am J Cardiol* 1996 Oct 15; 78:890-5.
- [13] Rogers WJ, Johnstone DE, Yusuf S, Weiner DH, Gallagher P, Bittner VA, et al. Quality of life among 5,025 patients with left ventricular dysfunction randomized between placebo and enalapril: the Studies of Left Ventricular Dysfunction. The SOLVD Investigators. *J Am Coll Cardiol* 1994 Feb; 23:393-400.
- [14] Evangelista LS, Berg J, Dracup K. Relationship between psychosocial variables and compliance in patients with heart failure. *Heart Lung* 2001 Jul; 30:294-301.
- [15] Nieuwenhuis MM, Jaarsma T, van Veldhuisen DJ, van der Wal MH. Self-reported versus 'true' adherence in heart failure patients: a study using the Medication Event Monitoring System. *Neth Heart J* 2012 Aug; 20 (7-8):313-9.
- [16] Van der Wal MH, Jaarsma T, Moser DK, Veeger NJ, van Gilst WH, van Veldhuisen DJ. Compliance in heart failure patients: the importance of knowledge and beliefs. *Eur Heart J* 2006 Feb; 27:434-40.
- [17] Evangelista LS, Berg J, Dracup K. Relationship between psychosocial variables and compliance in patients with heart failure. *Heart Lung* 2001 Jul; 30:294-301.
- [18] Nieuwenhuis MM, Jaarsma T, van Veldhuisen DJ, van der Wal MH. Self-reported versus 'true' adherence in heart failure patients: a study using the Medication Event Monitoring System. *Neth Heart J* 2012 Aug; 20 (7-8):313-9.
- [19] Van der Wal MH, van Veldhuisen DJ, Veeger NJ, Rutten FH, Jaarsma T. Compliance with non-pharmacological recommendations and outcome in heart failure patients. *Eur Heart J* 2010 Jun; 31:1486-93.
- [20] Van der Wal MH, Jaarsma T, van Veldhuisen DJ. Non-compliance in patients with heart failure; how can we manage it? *Eur J Heart Fail* 2005 Jan; 7:5-17.
- [21] Morcillo C, Aguado O, Delas J, Rosell F. [Utility of the Minnesota Living With Heart Failure Questionnaire for assessing quality of life in heart failure patients]. *Rev Esp Cardiol* 2007 Oct; 60:1093-6.
- [22] Kozhakenova LG, Lanzoni M, Rakhypbekov TK, Mussakhanova AK, Zurikanov KS, Castaldi S. Health-related Quality of Life in Kazakh Heart Failure Patients evaluated by the Minnesota Living with Heart Failure Questionnaire and comparison with a published large international sample. *Ann Ig* 2014 Nov; 26:547-52.
- [23] Lambrinou E, Kalogirou F, Lamnisis D, Middleton N, Sourtzi P, Lemonidou C, et al. Evaluation of the psychometric properties of the Greek version of the Minnesota Living With Heart Failure questionnaire. *J Cardiopulm Rehabil Prev* 2013 Jul; 33:229-33.
- [24] Naveiro-Rilo JC, Diez-Juarez DM, Romero BA, Rebollo-Gutierrez F, Rodriguez-Martinez A, Rodriguez-Garcia MA. Validation of the Minnesota living with heart failure questionnaire in primary care. *Rev Esp Cardiol* 2010 Dec; 63:1419-27.
- [25] Parajon T, Lupon J, Gonzalez B, Urrutia A, Altimir S, Coll R, et al. [Use of the Minnesota Living With Heart Failure Quality of Life Questionnaire in Spain]. *Rev Esp Cardiol* 2004 Feb; 57:155-60.
- [26] Baghianimoghdam MH, Shogafard G, Sanati HR, Baghianimoghdam B, Mazloomi SS, Askarshahi M. Application of the health belief model in promotion of self-care in heart failure patients. *Acta Med Iran* 2013; 51:52-8.
- [27] Yayehd K, Damorou F, N'cho Mottoh MP, Tcherou T, Johnson A, Pessinaba S, et al. [Compliance to treatment in heart failure patients in Lome]. *Ann Cardiol Angeiol (Paris)* 2013 Feb; 62:22-7.
- [28] Kato N, Kinugawa K, Seki S, Shiga T, Hatano M, Yao A, et al. Quality of life as an independent predictor for cardiac events and death in patients with heart failure. *Circ J* 2011; 75:1661-9.
- [29] Heo S, Moser DK, Lennie TA, Zambroski CH, Chung ML. A comparison of health-related quality of life between older adults with heart failure and healthy older adults. *Heart Lung* 2007 Jan; 36:16-24.
- [30] Gott M, Barnes S, Parker C, Payne S, Seamark D, Gariballa S, et al. Predictors of the quality of life of older people with heart failure recruited from primary care. *Age Ageing* 2006 Mar; 35:172-7.
- [31] Lewis EF, Lamas GA, O'Meara E, Granger CB, Dunlap ME, McKelvie RS, et al. Characterization of health-related quality of life in heart failure patients with preserved versus low ejection fraction in CHARM. *Eur J Heart Fail* 2007 Jan; 9:83-91.
- [32] Luttik ML, Jaarsma T, Veeger N, Van Veldhuisen DJ. Marital status, quality of life, and clinical outcome in patients with heart failure. *Heart Lung* 2006 Jan; 35:3-8.