
Analysis of Cosmic Rays from Inter-Electronic Structure of the Electron-Revised Version

D. P. Nandedkar*

Department of Electrical Engineering, Indian Institute of Technology, Bombay, Powai, Mumbai, India

Abstract

Here an analysis of cosmic rays from inter-electronic structure of the electron is reconsidered with respect to a previous paper by Nandedkar, where electromagnetic nature of cosmic rays is analyzed along with complex charge and complex mass of an electron. Here it is interesting to note that the moving mass of the electron inside and / on periphery of the electron has a velocity of $4.239(4) \times 10^8$ m/ sc which is greater than velocity of light in free space. Thus a world of Tachyon (where particle velocity can exceed velocity of light in free space) exit inside and / on periphery of the electron. In this paper strength of magnetic flux density required for reversal of spin angular momentum is reconsidered with real part of complex charge, of the moving electron along periphery of the circle of electron having radius of $1.878(8) \times 10^{-15}$ m, illustrating spin of the electron diagrammatically. This electron radius of $1.878(8) \times 10^{-15}$ m called “reduced radius” is (2/3) times so called “classical radius” of electron. Distribution of complex charge and complex mass of the electron is confined to the circle of “reduced radius” of the electron. Strength of magnetic flux density required for reversal of spin angular momentum is 1.877×10^{14} Wb / m². Sudden alternate reversals of magnetic flux densities in this case of equal to or greater than above value, generate continuous waves of cosmic rays of frequency $\sim 1.795(6) \times 10^{22}$ Hz by complex mass/ complex charge inter-electronic structure, which is reconsidered.

Keywords

Cosmic-Rays, Electron, Complex-Mass, Complex-Charge

Received: June 3, 2016 / Accepted: June 21, 2016 / Published online: August 5, 2016

@ 2016 The Authors. Published by American Institute of Science. This Open Access article is under the CC BY license.

<http://creativecommons.org/licenses/by/4.0/>

1. Introduction

Cosmic rays are extremely penetrating radiations/ charged particles, coming from where we do not know for certain as yet, but doubtless from regions far away from earth, and continually bombarding it on all sides with more or less of uniform intensity.

It is considered that the cosmic rays have terrestrial origin. As regards nature of charged particles of which cosmic rays are composed of, whether positive or negative, protons or electrons, study of the latitude effect on earth, by itself, gives no indication whatever; nor does it excludes the possibility of the presence of photons or neutrons, [1-7].

In a previous paper by Nandedkar [8], an analysis of cosmic rays from inter-electronic structure of the electron is considered, where electromagnetic nature of cosmic rays is analyzed along with complex charge and complex mass of an electron. Here it is interesting to note that the moving mass of the electron inside and / on periphery of the electron has a velocity of $4.239(4) \times 10^8$ m / sc which is greater than velocity of light in free space. Thus a world of Tachyon [9] (where particle velocity can exceed velocity of light in free space) exit inside and/ on periphery of the electron. In this paper strength of magnetic flux density required for reversal

* Corresponding author

E-mail address: dpn@ee.iitb.ac.in

of spin angular momentum is reconsidered with real part of complex charge, of the moving electron along periphery of the circle of electron having radius of $1.878(8) \times 10^{-15}$ m, illustrating spin of the electron diagrammatically. This electron radius of $1.878(8) \times 10^{-15}$ m called “reduced radius” is (2/3) times so called “classical radius” of electron. Distribution of complex charge and complex mass of the electron is confined to the circle of “reduced radius” of the electron. Strength of magnetic flux density required for reversal of spin angular momentum is $1.877(1) \times 10^{14}$ Wb/ m². Sudden alternate reversals of magnetic flux densities in this case of equal to or greater than above value, generate continuous waves of cosmic rays of frequency $\sim 1.795(6) \times 10^{22}$ Hz by complex mass/ complex charge inter-electronic structure, which is reconsidered.

Frequencies higher than about 10^{21} Hz of Electromagnetic Waves are classified in Cosmic Rays. In this research-paper, aspect of electromagnetic nature of the cosmic rays is reconsidered in detail incorporating some additional data with reference to previous paper by Nandedkar [8].

Figures in this article are in, Plane of Paper for hard-copy version/ Plane of Flat Monitor of Computer for soft-copy version, unless otherwise specified.

The article is developed in following Sections:

1. Introduction
2. Radius of an Electron
3. Complex Charge and Complex mass of an Electron
4. Electromagnetic Radiation Generation by the Electron
5. Numerical Analysis and
6. Conclusions

2. Radius of an Electron

Let a small uniformly charged sphere O i.e. the electron of radius r_e and charge e moves along the X-axis with a steady velocity u . If this velocity is not large, it may be assumed that the sphere carries its Faradays tubes along with it, undisturbed as the electromagnetic induction that tends to distort the tubes of force, depends on the velocity of motion - (Fig 1).

As the moving charge O is equivalent to a current i of strength $i = eu$, the magnetic field H due to it at a point P, distant r from O is given by Biot (Ampere) Law as follows,

$$H = \frac{eu \sin \theta}{4\pi r^2}, \tag{1}$$

where θ is the angle between r and X-axis. Therefore energy density w of magnetic field [10] at P, is given by,

$$w = \frac{1}{2} \mu_0 H^2. \tag{2}$$

where μ_0 is permeability of free space.

Now consider a small element PQRS at P (Fig. 1) with $PQ = dr$ and PS subtending at O an angle $d\theta$, so that $PS = r d\theta$. The area of the element PQRS (in the plane of the Fig. 1) = $r d\theta \times dr$. Dropping a perpendicular $PN = r \sin \theta$ on the X-axis from P, if the small area PQRS considered above at P, be revolved about OX-axis then it will enclose a ring, every bit of which lies upon the circumference of the circle whose radius is $PN = r \sin \theta$ and whose plane is perpendicular to OX and hence has the same value for H .




Fig. 1. Diagram for considering of magnetic field energy of a moving electron of finite size with velocity u along X-axis.

The volume of this ring is dv' , where

$$dv' = (rd\theta \times dr) [2\pi \times (PN = r \sin \theta)]$$

which gives,

$$dv' = 2\pi r^2 \sin \theta d\theta dr. \tag{3}$$

The contribution to the differential energy of the magnetic field dw at dv' is given, using eqns. (1), (2) and (3), by

$$dw = w dv' = \frac{\mu_0}{16\pi} \frac{e^2 u^2 \sin^3 \theta}{r^2} d\theta dr. \tag{4}$$

Integrating (Fig. 1) the above quantity given by eqn. (4), for all values r i.e. from $OP = r_e$ to ∞ (where r_e is the radius of electron), and for all values of θ , i.e. from 0 to π , total energy due to the moving charged sphere i.e. the electron, is given by,

$$W = \frac{\mu_0}{16\pi} e^2 u^2 \int_0^\pi \sin^3 \theta d\theta \int_{r_e}^\infty \frac{dr}{r^2} = \frac{\mu_0 e^2 u^2}{12 \pi r_e}. \tag{5}$$

The above energy, is also the kinetic energy E of electron, given by,

$$E = \frac{1}{2} m_0 u^2, \tag{6}$$

here m_0 is rest mass of electron, assuming,

$$u \ll c, \tag{7}$$

where c is the velocity of an electromagnetic wave in free space.

Equating eqns. (5) and (6) gives,

$$m_o = \frac{\mu_o e^2}{6 \pi r_e}, \tag{8}$$

here the mass m_o is due to motion where eqn. (7) holds good and, energy associated with the field is given by eqn. (6), in Newtonian World.

Now [10],

$$c = \frac{1}{\sqrt{\mu_o \epsilon_o}}, \tag{9}$$

so,

$$r_e = \frac{(\frac{2}{3})e^2}{4 \pi \epsilon_o m_o c^2}, \tag{10}$$

using eqns. (8) and (9). Eqn. (10) gives radius of an electron. Here ϵ_o is permittivity of free space. Now classical radius of electron r_{ec} is given [11] by,

$$r_{ec} = \frac{e^2}{4 \pi \epsilon_o m_o c^2}. \tag{11}$$

From eqns. (10) and (11),

$$r_e / r_{ec} = 2/3. \tag{12}$$

In view of eqn. (12), r_e of eqn. (10) is called “reduced radius” of the electron, whereas eqn. (11) gives “classical radius” of the electron. In this analysis “radius” word is used for “reduced radius” of the electron, unless otherwise specified.

3. Complex Charge and Complex Mass of an Electron

From eqn. (10),

$$m_o c^2 = \frac{(\frac{2}{3})e^2}{4 \pi \epsilon_o r_e}. \tag{13}$$

In eqn. (13), $m_o c^2$ is the total energy associated with rest mass m_o of the electron [12]. Here $\frac{(\frac{2}{3})e^2}{4 \pi \epsilon_o r_e}$ gives Potential Energy of either charges e_1 or e_2 due to e_2 or e_1 separated by a distance r_e - (here also refer to Secn. 4), such that,

$$e_1 + e_2 = e, \tag{14}$$

and,

$$e_1 e_2 = (\frac{2}{3}) e^2. \tag{15}$$

The solution of simultaneous equations denoted by eqns. (14) and (15) in e_1 and e_2 , gives that,

$$e_1 = \frac{e}{2} (1 + i\sqrt{5/3}), \tag{16}$$

and,

$$e_2 = \frac{e}{2} (1 - i\sqrt{5/3}), \tag{17}$$

where $i = \sqrt{-1}$.

Let $(m_o - im'_o)$ be the rest mass of charge e_1 and im'_o be the rest mass of charge e_2 , such that total rest mass of charges e_1 and e_2 is $(m_o - im'_o) + im'_o = m_o$ and total rest mass energy of them is,

$$[(m_o - im'_o) + im'_o]c^2 = m_o c^2. \tag{18}$$

And then eqn. (13) can be rewritten as follows:

$$\frac{(\frac{e}{2}(1+i\sqrt{5/3}))\{(\frac{e}{2}(1-i\sqrt{5/3}))\}}{4 \pi \epsilon_o r_e} = \{(m_o - im'_o) + im'_o\} c^2, \tag{19}$$

where, $(m_o - im'_o)$ is rest mass of charge $e_1 = \frac{e}{2} (1 + i\sqrt{5/3})$, and im'_o is rest mass of charge $e_2 = \frac{e}{2} (1 - i\sqrt{5/3})$, using eqns. (15), (16), (17) and (18).

4. Electromagnetic Radiation Generation by the Electron




Fig. 2. Complex charge and complex mass structure of electron of radius r_e in (p, ϕ) -plane of a system of polar co-ordinates. Here imaginary-axis is not shown in above Figure. Entities associated with a factor $i = \sqrt{-1}$ are mathematical operators only.

In this model of electron, the charge e_1 with rest mass $(m_o - im'_o)$ is present at origin O of a system of polar co-ordinates (p, ϕ) . And charge e_2 rotate along circle of radius r_e with charge e_1 hinged at O - (Fig. 2), forming a stable system, is the assumption. The plane of Fig. 2 is Plane of Paper.

Now potential energy of charge e_2 on circumference of circle of radius r_e due to potential of charge e_1 at O (Fig. 2) is

$[e_2 (\frac{e_1}{4 \pi \epsilon_0 r_e})] = \frac{(\frac{2}{3})e^2}{4 \pi \epsilon_0 r_e} = m_0 c^2$ [refer to eqns. (13) and (15)]. Similarly potential energy of charge e_1 at center O due to potential of charge e_2 on circumference of circle of radius r_e (Fig. 2) is $[e_1 (\frac{e_2}{4 \pi \epsilon_0 r_e})] = \frac{(\frac{2}{3})e^2}{4 \pi \epsilon_0 r_e} = m_0 c^2$ [refer to eqns. (13) and (15)]. So this rest mass energy of the electron $m_0 c^2$ is also effective pair potential energy that describes the interaction which acts along the line of length equal to the radius r_e of electron connecting the (two point) charges e_1 and e_2 .

Let m'_{oe} be the moving mass of charge e_2 . Now *the assumption* is that, this charge of moving mass moves with uniform velocity v along the circumference of the circle of radius r_e (Fig. 2). Then,

$$m'_{oe} = \frac{im'_0}{\sqrt{1-\frac{v^2}{c^2}}}, \quad (20)$$

by Einstein theory of relativity [12].

If I is the moment of Inertia of the spinning electron about O, then

$$I = m'_{oe} r_e^2. \quad (21)$$

And spinning energy associated with this charge e_2 is then,

$$E_s = \frac{1}{2} I \omega^2 = \frac{1}{2} m'_{oe} r_e^2 \omega^2, \quad (22)$$

where ω is the angular velocity of the spinning/rotating electron in the circle of radius r_e .

Further energy associated with moving charge of mass m'_{oe} , by Einstein theory of relativity [12], is given as follows:

$$E_m = m'_{oe} c^2. \quad (23)$$

Here E_s of eqn. (22) and, E_m of eqn. (23) are same. Hence equating the two, the result is.

$$\omega = \sqrt{2} \frac{c}{r_e}. \quad (24)$$

But spin angular momentum of charge e_2 is,

$$M_s = I \omega = m'_{oe} r_e^2 \omega, \quad (25)$$

using eqn. (21).

Further, the spin angular momentum of the electron is considered to be given by,

$$M_s = \pm \frac{1}{2} \left(\frac{h}{2\pi} \right), \quad (26)$$

where, $\pm 1/2$ is spin quantum number, say $+1/2$ for clockwise spin & $-1/2$ for anti-clockwise spin of the electron and, h is Plank constant, in analogy with Uhlenbeck and Goudschmidt (1925-1926) model of spinning electron [13]. Eqn. (26) is *the*

assumption of this analysis.

From eqn. (26) choosing $+1/2$ spin quantum number, eqn. (25) gives,

$$m'_{oe} = \frac{1}{2} \left(\frac{h}{2\pi} \right) \frac{1}{r_e^2 \omega}. \quad (27)$$

Substituting the value of ω from eqn. (24) in eqn. (27), eqn. (27) gives,

$$m'_{oe} = \frac{1}{2\sqrt{2}} \left(\frac{h}{c r_e} \right), \quad (28)$$

where,

$$\hbar = h/2\pi, \quad (29)$$

is reduced Planck constant.

Further, since

$$m'_{oe} = \frac{im'_0}{\sqrt{1-\frac{v^2}{c^2}}},$$

which is eqn. (20) and,

$$r_e \omega = v = \sqrt{2} c, \quad (30)$$

using eqn. (24), hence eqn. (20) using eqn. (30), gives,

$$m'_{oe} = m'_0 = \frac{1}{2\sqrt{2}} \left(\frac{h}{c r_e} \right), \quad (31)$$

using eqn. (28).

Energy associated with spin angular momentum $+(1/2) \hbar$ [refer to eqn. (26) and eqn. (29)] is $m'_{oe} c^2$, say in clockwise spin of the electron [refer to eqn. (23)]. Similarly, energy associated with spin angular momentum $-(1/2) \hbar$ [refer to eqn. (26) and eqn. (29)] is $-m'_{oe} c^2$, say in anticlockwise spin of the electron [refer to eqn. (23)]. So change of energy of the electron when its spin momentum changes from $+(1/2) \hbar$ to $-(1/2) \hbar$ is given by,

$$[m'_{oe} c^2 - (-m'_{oe} c^2)] = 2 m'_{oe} c^2. \quad (32)$$

The above difference of energy of the electron is radiated with a photon of frequency ν of energy $h\nu$, by Planck hypothesis. Whence Eqn. (32) gives,

$$h\nu = 2 m'_{oe} c^2. \quad (33)$$

Using eqn. (31), eqn. (33) gives,

$$\nu = \frac{1}{2\pi\sqrt{2}} \left(\frac{c}{r_e} \right). \quad (34)$$

Equation (34) gives frequency of electromagnetic radiation from the electron as its spin angular momentum changes from $+(1/2) \hbar$ to $-(1/2) \hbar$.

5. Numerical Analysis

Here numerical values of various entities with reference to previous paper by Nandedkar [8] are mentioned below (for ready reference), using “Data / Table 16/ General Physical Constants” of Appendix 9, pp. 1003 [14]:

[1]. Radius of the electron (r_e):

$$r_e = \frac{\left(\frac{2}{3}\right)e^2}{4\pi\epsilon_0 m_0 c^2} = 1.878(8) \times 10^{-15} \text{ m}, \quad (35)$$

refer to eqn. (10).

[2]. Velocity of the electron (v):

$$v = \sqrt{2} c = 4.239(4) \times 10^8 \text{ m/ sc}, \quad (36)$$

refer to eqn. (30).

This velocity is greater than velocity of electromagnetic radiation in free space, which can happen in Tachyon world [9].

[3]. Angular velocity of the electron (ω):

$$\omega = \sqrt{2} c / r_e = 2.256(4) \times 10^{23} \text{ rad/ sc}, \quad (37)$$

refer to eqn. (30).

[4]. Moving mass of the electron (m'_{oe}):

$$m'_{oe} = \frac{1}{2\sqrt{2}} \left(\frac{\hbar}{c r_e} \right) = 6.618(0) \times 10^{-29} \text{ Kgm}, \quad (38)$$

refer to eqn. (31).

[5]. Frequency of electromagnetic radiation from the electron (ν):

$$\nu = \frac{1}{2\pi\sqrt{2}} \left(\frac{c}{r_e} \right) = 1.795(6) \times 10^{22} \text{ Hz} \quad (39)$$

[refer to eqn. (34)], which is due to reversal of spin momentum of the electron. ν is in the range of cosmic rays.

[6]. Wavelength of electromagnetic radiation from the electron (λ):

$$\lambda = \frac{c}{\nu} = 2\pi\sqrt{2} r_e = 1.669(5) \times 10^{-14} \text{ m}, \quad (40)$$

[refer to eqn. (39)], which is due to reversal of spin momentum of the electron. λ is in the range of cosmic rays.

[7]. Strength of magnetic field required for reversal of spin angular momentum (B):

Here, value of magnetic flux density due to circulation of complex electronic current giving Real part of current due to Real charge along positive Y-axis (Fig. 3) is carried out in following steps.

Now magnetic flux density B [where instantaneous position of the electron denoted by arbitrary point Q coincides with point A on the circular electronic path shown in Fig. 3] developed

normal to and entering along negative Z -axis towards O and in the XY -plane of circular motion of the moving (say, clockwise) real charge $\frac{e}{2}$ of Complex Charge (moving), $e_2 = \frac{e}{2} (1 - i\sqrt{5/3})$ of eqn. (17) with velocity v of the electron which is in XY -plane of the paper, of the Fig. 3 produces Lorentz Force F_B (assumed to hold good) acting radially towards Centre O (Fig. 3) and along negative X -axis in XY -plane of paper of the Fig. 3, that is given by following relationship:

$$F_B = B \frac{e}{2} v. \quad (41)$$




Fig. 3. Charge $e_2 = (e/2) (1 - i\sqrt{5/3})$ of electron rotating clockwise in a circle of electron radius r_e with moving mass m'_{oe} and velocity v in XY -plane, where imaginary-axis is not shown in above Figure. Entities associated with a factor $i = \sqrt{-1}$ are mathematical operators only. Charge $e_1 = (e/2) (1 + i\sqrt{5/3})$ of electron with its rest mass $(m_0 - im'_0)$ is at Center O of the circle of electron radius r_e . Here B = magnetic flux density developed (due to circulation of Real part of current due to Real charge along positive Y -axis is assumption, acting along negative Z -axis towards O , F_B = Lorentz Force acting towards O along negative X -axis in XY -plane, $-F_{Coul}$ = Coulomb Force of Repulsion acting towards O along negative X -axis in XY -plane, and F_C = Centrifugal Force acting away from O along positive X -axis in XY -plane. Orientation of X -axis in XY -plane is to be chosen such that v is always instantaneously tangential to path of circular motion of electron in XY -plane (say at A) and OX is normal to v along the tangent TA at A . Here Q denotes an arbitrary point on circular orbit of the electron of Fig. 3. Origin O of the system of Polar co-ordinates (p, ϕ) with (p, ϕ) - plane of Fig. 2, is same as origin O of the system of Cartesian co-ordinates (X, Y) with (X, Y) - plane of Fig. 3. Plane of Fig. 2 is same as plane of Fig. 3, which is the plane of paper. Radius of electron r_e is same in circles of the radius r_e (Fig. 2 & Fig. 3) and Center O of Fig. 2 coincides with Center O of Fig. 3, although both Figures are drawn with different scales.

Whereas Coulomb Force F_{Coul} (assumed to hold good) of repulsion between Charge (stationary), $e_1 = \frac{e}{2} (1 + i\sqrt{5/3})$ of eqn. (16) and Charge (moving), $e_2 = \frac{e}{2} (1 - i\sqrt{5/3})$ of eqn. (17), separated by distance of r_e of electron radius of Fig. 3 [where instantaneous position of the electron denoted by arbitrary point Q coincides with point A on the circular electronic path shown in Fig. 3 as mentioned before], is given by Coulomb Law, viz.,

$$F_{\text{Coul}} = \left(\frac{2}{3}\right) e^2/4 \pi \epsilon_0 r_e^2, \quad (42)$$

where,

$$\left\{\frac{e}{2} \left(1 + i \sqrt{\frac{5}{3}}\right)\right\} \left\{\frac{e}{2} \left(1 - i \sqrt{\frac{5}{3}}\right)\right\} = \left(\frac{2}{3}\right) e^2, \quad (43)$$

and where e_1 and e_2 are given by eqns. (16) and (17) respectively, [here refer also to eqn. (15)]. And here r_e is given by eqn. (10). $-F_{\text{Coul}}$ acts radially towards Centre O along negative X-axis in XY-plane of the paper of Fig. 3.

Thus a force due to difference of Lorentz and Coulomb forces, viz.,

$$F_B - F_{\text{Coul}}, \quad (44)$$

acts towards Center O of circular path of electron of Fig. 3 along negative X-axis in XY-plane of the paper.

Further Centrifugal Force F_C (*assumed to hold good*) acting on the above electron in the XY-plane of the paper [where instantaneous position of the electron denoted by arbitrary point Q coincides with point A on the circular electronic path shown in Fig. 3 as mentioned before], and radially away from Centre O along positive X-axis due to circular motion of the electron, is given by,

$$F_C = \left(\frac{m'_{oe} v^2}{r_e}\right), \quad (45)$$

where m'_{oe} , v and r_e are given by eqns. (28), (30) and (10) respectively.

Expressions given by eqns. (44) and (45) are the same. Whence equating them gives,

$$F_B - F_{\text{Coul}} = F_C. \quad (46)$$

Eqn. (46) using eqns. (41), (42) and (45), gives,

$$B = \left(\frac{\left(\frac{m'_{oe} v^2}{r_e} + \left(\frac{2}{3}\right) e^2/4 \pi \epsilon_0 r_e^2\right)}{ev/2}\right) = 1.877(1) \times 10^{14} \text{ Wb/m}^2. \quad (47)$$

6. Conclusions

An analysis of cosmic rays from inter-electronic structure of the electron is reconsidered with respect to a previous paper by Nandedkar [8], where electromagnetic nature of cosmic rays is analyzed along with complex charge and complex mass of an electron. Here it is interesting to note that the moving mass $6.618(0) \times 10^{-29}$ Kgm of the electron [eqn. (38)] inside and/ on periphery of the electron has a velocity of $4.239(4) \times 10^8$ m/ sc [eqn. (36)] which is greater than velocity of light in free space. Thus a world of Tachyon [9] (where particle velocity can exceed velocity of light in free space) exist inside and/ on periphery of the electron. In this

paper strength of magnetic flux density required for reversal of spin angular momentum is reconsidered with real part of complex charge, of the moving electron along periphery of the circle of electron having radius of $1.878(8) \times 10^{-15}$ m [(eqn. (35)], illustrating spin of the electron diagrammatically (Fig. 3). This electron radius of $1.878(8) \times 10^{-15}$ m called “reduced radius” is (2/3) times so called “classical radius” of electron [refer to eqns. (10, 11 & 12]. Distribution of complex charge and complex mass of the electron is confined to the circle of “reduced radius” of the electron (Fig. 2). Strength of magnetic flux density required for reversal of spin angular momentum is $1.877(1) \times 10^{14}$ Wb/ m² [eqn. (47)], as per (*assumed*) quantum condition of spin angular momentum $M_s = \pm \frac{1}{2} \left(\frac{h}{2\pi}\right)$ of eqn. (26). Sudden alternate reversals of magnetic flux densities in this case of equal to or greater than above value, generate continuous waves of cosmic rays of frequency $\sim 1.795(6) \times 10^{22}$ Hz [eqn. (39)] by complex mass/ complex charge inter-electronic structure reconsidered herewith.

Here note that potential energy of charge e_2 on circumference of circle of radius r_e due to potential of charge e_1 at O (Fig. 2) is $\left[e_2 \left(\frac{e_1}{4 \pi \epsilon_0 r_e}\right)\right] = \frac{\left(\frac{2}{3}\right)e^2}{4 \pi \epsilon_0 r_e} = m_0 c^2$ [refer to eqns. (13) and (15)]. Similarly potential energy of charge e_1 at center O due to potential of charge e_2 on circumference of circle of radius r_e (Fig. 2) is $\left[e_1 \left(\frac{e_2}{4 \pi \epsilon_0 r_e}\right)\right] = \frac{\left(\frac{2}{3}\right)e^2}{4 \pi \epsilon_0 r_e} = m_0 c^2$ [refer to eqns. (13) and (15)]. So this rest mass energy of the electron $m_0 c^2$ is also effective pair potential energy that describes the interaction which acts along the line of length equal to the radius r_e of electron connecting the (two point) charges e_1 and e_2 .

This research-paper illustrates diagrammatically spin of electron [Fig. 2 and Fig. 3 which are assumed to hold good. Here *the assumption* is that, the charge of moving mass m'_{oe} {eqn. (31)} moves with uniform velocity v {eqn. (30)} along the circumference of the circle of radius r_e {Fig. 2 and Fig. 3}}] with all entities associated therein with a factor $i = \sqrt{-1}$ as mathematical operators only, in Tachyon world [9] for the electron of charge e and mass m_0 of Newtonian world.

In this analysis Lorentz Force [eqn. (41)], Centrifugal Force [eqn. (45)] and Coulomb Force of Repulsion [eqn. (42)], are all assumed to hold good and are all Real. Balancing for moving electron (Fig. 3) under these Forces for *inward* Lorentz Force *towards Center O of circle* (Fig. 3) against *outward* Centrifugal Force & Coulomb Force of Repulsion with reference to stationary charge at Center O of circle of ‘reduced radius’ of electron *away from Center O of circle* (Fig. 3) considering say, instantaneous position of the electron denoted by arbitrary point Q coinciding with point A

on the circular electronic path shown in Fig. 3, gives magnetic flux density as Real & positive required for reversal of spin angular momentum as given by eqn. (47) of value $1.877(1) \times 10^{14} \text{ Wb/m}^2$.

If different species of electrons with different charge/ mass ratios and hence different radii be considered to exist in this vast universe, then a continuous/ discrete band of cosmic rays of various frequencies may be generated, depending upon what different species of electrons with different charge/ mass ratios and hence different radii are available in this vast universe. Data on energy spectra of cosmic ray (photons) at Cosmic Ray Observatory corresponding to Wikipedia, the free encyclopaedia, can also be considered to be due to production of cosmic rays mentioned in this article.

Acknowledgements

The author is thankful for the interesting discussions with reference to this research-paper given herewith he had, with his Guru (the Teacher) Dr. Ing. G. K. Bhagavat, [(now) Late] Professor of Electrical Engineering Department, Indian Institute of Technology (IIT), Bombay, Powai, Mumbai, India.

Further he gives thanks also to Electrical Engineering Department, IIT, Mumbai, India for giving permission to publish this research-work.

References

- [1] Michael A. DuVernois (Editor) (1999): "Topics in Cosmic-Ray Astrophysics", Nova Science Publishers Inc, USA.
- [2] Ringuet L. L., (1950): "Cosmic Rays", Prentice-Hall, New York.
- [3] A. Dauvillier, (1954): "Les Rayons Cosmiques", Dunod publisher, Paris, France.
- [4] Wilson J. G.-Editor, (1952): "Progress in cosmic ray physics", Vol. I, North Holland, Amsterdam.
- [5] Wilson J. G.-Editor, (1954): "Progress in cosmic ray physics", Vol. II, North Holland, Amsterdam.
- [6] Wilson J. G.-Editor, (1956): "Progress in cosmic ray physics", Vol. III, North Holland, Amsterdam.
- [7] Mewaldt, R. A., (1996): "Comic Rays", Macmillan Encyclopaedia of Physics.
- [8] Nandedkar, D. P. (2016): "Analysis of cosmic rays from inter-electronic structure of the electron", Vol. 2, No. 3, pp. 220-225, Phys. Journal (PSF), (AIS), USA.
- [9] Feinberg, G. (1967). "Possibility of Faster-Than-Light Particles". Physical Review 159 (5): 1089–1105.
- [10] Ramo, S., Whinnery, J. R. and Van Duzer, T., (1970): "Fields and Waves in Communication Electronics", Wiley Eastern Private Limited", First Edition, First Wiley Eastern Reprint, New Delhi.
- [11] French, A. P., (1958): "Principles of Modern Physics", Wiley International Edition, New York, London etc.
- [12] Richtmyer, F. K., Kennard, E. H., and Lauritsen, T., (1955): "Introduction to Modern Physics", Fifth Edition, International Student Edition, McGraw Hill Book Co, Inc., New York, Toronto, London etc.
- [13] Jain, G. C., (1967): "Properties of Electrical Engineering Materials", Harper & Row, Publishers, New York, Evanston and London.
- [14] Parker, P., (1957): "Electronics", Reprinted Published First Edition, Edward Arnold Publishers Ltd., London.