
Solution of Eighth Order Boundary Value Problem by Using Variational Iteration Method

Abdulla-Al-Mamun^{*}, Mohammad Asaduzzaman, Samsun Nahar Ananna

Department of Mathematics, Islamic University, Kushtia, Bangladesh

Abstract

In this paper, we introduce some basic idea of Variational iteration method for short (VIM) to solve the eighth order boundary value problems. It is to be mentioned that, presently, the literature on the numerical solutions of eighth order boundary value problem and associated eigen value problems is not available. By using a suitable transformation, the variational iteration method can be used to show that eighth order boundary value problems are equivalent to a system of integral equation. The VIM is used to solve effectively, easily, and accurately a large class of non-linear problems with approximations which converge rapidly to accurate solutions. For linear problems, it's exact solution can be obtained by only one iteration step due to the fact that the Lagrange multiplier can be exactly identified. It is to be noted that the Lagrange multiplier reduces the iteration on integral operator and also minimizes the computational time. The method requires no transformation or linearization of any forms. Two numerical examples are presented to show the effectiveness and efficiency of the method. Also, we compare the result with exact solution. Finally, we investigate the error between numerical solution and exact solution and draw the graph of error function by using Mathematica.

Keywords

Variational Iteration Method, Boundary Value Problem, Mathematica, Exact Solution, Approximate Solution, Numerical Solution, Lagrange Multiplier, Absolute Error

Received: March 29, 2019 / Accepted: May 14, 2019 / Published online: May 27, 2019

© 2019 The Authors. Published by American Institute of Science. This Open Access article is under the CC BY license.

<http://creativecommons.org/licenses/by/4.0/>

1. Introduction

Consider the general eighth order boundary value problem of the type:

$$\left. \begin{aligned} u^{(8)} &= f(x, u(x)), a \leq x \leq b \\ u^{(i)}(a) &= A_i, i = 0,1,2,3 \\ u^{(j)}(b) &= B_j, j = 0,1,2,3 \end{aligned} \right\} \quad (1)$$

Where $A_i, i = 0,1,2,3$ and $B_j, j = 0,1,2$ are finite real constants, also $f(x, u(x))$ is a continuous function on $[a, b]$.

Eighth order boundary value problems are known to arise in the mathematics, physics and engineering sciences [1]. In

addition, it is well known that when a layer of fluid is heated from below and is subject to the action of rotation, instability may set in as over stability [2, 3]. This instability may be modeled by an eighth order ordinary differential equation with appropriate boundary conditions [3, 4]. For more discussion about the eighth-order boundary value problems, see [2, 4-7] and the references therein. The literature of numerical analysis contains little on the solution of the eighth-order boundary value problems [5]. Research in this direction may be considered in its early stages. Theorems which list the conditions for the existence and uniqueness of solutions of such problems are contained in a comprehensive survey by Agarwal [8].

The boundary value problems of higher order have been

* Corresponding author

E-mail address: abdullamamun21@gmail.com (Abdulla-Al-Mamun)

investigated because of both of their mathematical importance and the potential for applications in hydrodynamic and hydromagnetic stability. Finite-difference method was employed in [6, 8] to find the solution of eighth-order boundary value problems. The obtained results were divergent at points adjacent to the boundary. In a later study, Siddiqi and Twizell [3] used octic polynomial spline for solving these problems. Twizell et al. [3, 5, 7] also solved some other higher-order problems and encountered the same deficiencies. The divergent results are due to the use of lower-order test function in the spline methods. The spline function values at the mid knots of the interpolation interval and the corresponding values of the even-order derivatives are related through consistency relations. We apply the variational iteration method (VIM) to find solutions of eighth-order boundary value problems.

2. History of Variation Method (VIM)

The variational iteration method (VIM) developed in 1999 by He in [9-19] will be used to study the linear wave equation, nonlinear wave equation, and wave-like equation in bounded and unbounded domains. The method has been proved by many authors [20-31] to be reliable and efficient for a wide variety of scientific applications, linear and nonlinear as well. It was shown by many authors that this method is more powerful than existing techniques such as the Adomian method [32, 33], perturbation method, etc. The method gives rapidly convergent successive approximations of the exact solution if such a solution exists; otherwise a few approximations can be used for numerical purposes. The method is effectively used in [10-25] and the references therein. The perturbation method suffers from the computational workload, especially when the degree of nonlinearity increases. Moreover, the Adomian method suffers from the complicated algorithms used to calculate the Adomian polynomials that are necessary for nonlinear problems. The VIM has no specific requirements, such as linearization, small parameters, etc. for nonlinear operators.

The variational iteration method, which is a modified general Lagrange multiplier method has been shown to solve effectively, easily and accurately, a large class of nonlinear problems with approximations which converge quickly to accurate solutions. It was successfully applied to autonomous ordinary differential equation [13], to nonlinear partial differential equations with variable coefficients [34], to

Schrödinger-KDV, generalized KDV and shallow water equations [35], to Burgers' and coupled Burgers' equations [36], to the linear Helmholtz partial differential equation [38] and recently to nonlinear fractional differential equations with Caputo differential derivative [40], and other fields [9, 10, 20, 39, 40-43]. On the other hand, one of the interesting topics among researchers is solving integro-differential equations.

3. Basic Ideas of Variational Iteration Method

To illustrate the basic concept and idea of the variational iteration technique, we consider the following general differential equation

$$Lf + Nf = g(x) \quad (2)$$

where L is a linear operator, N a nonlinear operator and g (x) is inhomogeneous forcing term. According to the variational iteration method [9], we can construct a correct functional as follows

$$f_{n+1}(x) = f_n(x) + \int_0^x \lambda (Lf_n(s) + L\tilde{f}_n(s) - g(s)) ds \quad (3)$$

Where λ is a Lagrange multiplier, which can be identified optimally via the variational iteration method. The subscripts n denoted the nth approximation, \tilde{u}_n is considered as a restricted variation. i.e. $\delta\tilde{f}_n = 0$; (3) is called as a correct functional.

The solution of linear problems can be solved in a single iteration step due to the exact identification of their Lagrange multiplier.

For the sake of simplicity, we consider the following system of differential equations:

$$x'_i(t) = u_i(t, x_i), i = 1, 2, 3, \dots, n \quad (4)$$

subject to the boundary conditions, $x_i(0) = c_i, i = 1, 2, 3, \dots, n$.

To solve the system by means of the variational iteration method, we can rewrite the system (4) in the following form:

$$x'_i(t) = u_i(x_i) + g_i(t), i = 1, 2, 3, \dots, n \quad (5)$$

subject to the boundary conditions, $x_i(0) = c_i, i = 1, 2, 3, \dots, n$ and g_i is defined in (2).

The correct functional for the nonlinear system (1.04) can be expressed as follows:

$$\left. \begin{aligned} x_1^{(k+1)}(t) &= x_1^{(k)}(t) + \int_0^t \lambda_1 \left(x_1'^{(k)}(T), f_1 \left(\widetilde{x}_1^{(k)}(T), \widetilde{x}_2^{(k)}(T), \dots, \widetilde{x}_n^{(k)}(T) \right) - g_1(T) \right) dT \\ x_2^{(k+1)}(t) &= x_2^{(k)}(t) + \int_0^t \lambda_2 \left(x_2'^{(k)}(T), f_2 \left(\widetilde{x}_1^{(k)}(T), \widetilde{x}_2^{(k)}(T), \dots, \widetilde{x}_n^{(k)}(T) \right) - g_2(T) \right) dT \\ &\dots \dots \dots \\ x_n^{(k+1)}(t) &= x_n^{(k)}(t) + \int_0^t \lambda_n \left(x_n'^{(k)}(T), f_n \left(\widetilde{x}_1^{(k)}(T), \widetilde{x}_2^{(k)}(T), \dots, \widetilde{x}_n^{(k)}(T) \right) - g_n(T) \right) dT \end{aligned} \right\} \quad (6)$$

where $\lambda_i, i = 1, 2, 3, \dots, n$ are Lagrange multipliers, $\widetilde{x}_1, \widetilde{x}_2, \dots, \widetilde{x}_n$ denote the restricted variations. Making the above functional stationary, we obtain the following conditions:

$$\begin{aligned} \lambda_i'(T)|_{T=t} &= 0, \\ 1 + \lambda_i'(T)|_{T=t} &= 0 \end{aligned}$$

For $i = 1, 2, 3, \dots, n$. Therefore, the Lagrange multipliers can be easily identified as:

$$\lambda_i = -1, i = 1, 2, 3, \dots, n \quad (7)$$

Substituting (7) into the correct functional (6), we have the following iteration formulae:

$$\begin{aligned} x_1^{(k+1)}(t) &= x_1^{(k)}(t) - \int_0^t \left(x_1'^{(k)}(T), f_1 \left(\widetilde{x}_1^{(k)}(T), \widetilde{x}_2^{(k)}(T), \dots, \widetilde{x}_n^{(k)}(T) \right) - g_1(T) \right) dT \\ x_2^{(k+1)}(t) &= x_2^{(k)}(t) - \int_0^t \left(x_2'^{(k)}(T), f_2 \left(\widetilde{x}_1^{(k)}(T), \widetilde{x}_2^{(k)}(T), \dots, \widetilde{x}_n^{(k)}(T) \right) - g_2(T) \right) dT \\ &\dots \dots \dots \\ x_n^{(k+1)}(t) &= x_n^{(k)}(t) - \int_0^t \left(x_n'^{(k)}(T), f_n \left(\widetilde{x}_1^{(k)}(T), \widetilde{x}_2^{(k)}(T), \dots, \widetilde{x}_n^{(k)}(T) \right) - g_n(T) \right) dT \end{aligned}$$

If we start the initial approximations $x_i(0) = c_i, i = 1, 2, 3, \dots, n$, then the approximation can be completely determined; finally, we approximate the solution.

$$x_i(t) = \lim_{k \rightarrow \infty} x_i^{(k)}(t)$$

by the nth term $x_i^{(n)}(t)$ for $i = 1, 2, 3, \dots, n$.

$$\left. \begin{aligned} \frac{df}{dx} &= l(x), \frac{dl}{dx} = m(x), \frac{dm}{dx} = n(x), \frac{dn}{dx} = p(x) \\ \frac{dp}{dx} &= q(x), \frac{dq}{dx} = r(x), \frac{dr}{dx} = s(x), \frac{ds}{dx} = e^{-x} f^2(x) \end{aligned} \right\} \quad (9)$$

With,

$$\begin{aligned} f(0) &= 1, l(0) = A, m(0) = 1, n(0) = B, p(0) = 1, q(0) \\ &= C, r(0) = 1, s(0) = D \end{aligned}$$

The system of differential equation (9) can be written in terms of the following system of integral equations with Lagrange multipliers $\lambda_i = +1; i = 1, 2, 3, \dots, 8$.

$$\begin{aligned} f^{(k+1)}(x) &= 1 + \int_0^x l^{(k)}(t) dt \\ l^{(k+1)}(x) &= A + \int_0^x m^{(k)}(t) dt \\ m^{(k+1)}(x) &= 1 + \int_0^x n^{(k)}(t) dt \\ n^{(k+1)}(x) &= B + \int_0^x p^{(k)}(t) dt \end{aligned}$$

4. Numerical Examples

In this paper, we present two examples to show efficiency and high accuracy of the variational iteration method for solving eighth order boundary value problems.

Example 01 Consider the following 8th order nonlinear boundary value problem

$$\left. \begin{aligned} f^{(8)}(x) &= e^{-x} f^2(x), 0 < x < 1 \\ f(0) &= f^{(2)}(0) = f^{(4)}(0) = f^{(6)}(0) = 1 \\ f(1) &= f^{(2)}(1) = f^{(4)}(1) = f^{(6)}(1) = e \end{aligned} \right\} \quad (8)$$

The exact solution of this problem is $f(x) = e^x$

The given 8th order boundary value problem can be transformed with the following system

$$p^{(k+1)}(x) = 1 + \int_0^x q^{(k)}(t) dt$$

$$q^{(k+1)}(x) = C + \int_0^x r^{(k)}(t) dt$$

$$r^{(k+1)}(x) = 1 + \int_0^x s^{(k)}(t) dt$$

$$s^{(k+1)}(x) = D + \int_0^x e^{-t} f^2(t) dt$$

With

$$f(0) = 1, l(0) = A, m(0) = 1, n(0) = B, p(0) = 1, q(0) = C, r(0) = 1, s(0) = D$$

Now, $f^{(1)}(x) = 1 + Ax$

$$l^{(1)}(x) = A + x$$

$$m^{(1)}(x) = 1 + Bx$$

$$n^{(1)}(x) = B + x$$

$$p^{(1)}(x) = 1 + Cx$$

$$q^{(1)}(x) = C + x$$

$$r^{(1)}(x) = 1 + Dx$$

$$s^{(1)}(x) = 1 + D - e^{-x}$$

$$f^{(2)}(x) = 1 + Ax + \frac{x^2}{2}$$

$$l^{(2)}(x) = A + x + \frac{B}{2}x^2$$

$$m^{(2)}(x) = 1 + Bx + \frac{x^2}{2}$$

$$n^{(2)}(x) = B + x + \frac{C}{2}x^2$$

$$p^{(2)}(x) = 1 + Cx + \frac{x^2}{2}$$

$$q^{(2)}(x) = C + x + \frac{D}{2}x^2$$

$$f^{(3)}(x) = 1 + Ax + \frac{x^2}{2} + \frac{B}{6}x^3$$

$$l^{(3)}(x) = A + x + \frac{B}{2}x^2 + \frac{1}{6}x^3$$

$$m^{(3)}(x) = 1 + Bx + \frac{1}{2}x^2 + \frac{C}{6}x^3$$

$$n^{(3)}(x) = B + x + \frac{C}{2}x^2 + \frac{1}{6}x^3$$

$$p^{(3)}(x) = 1 + Cx + \frac{1}{2}x^2 + \frac{D}{6}x^3$$

$$f^{(4)}(x) = 1 + Ax + \frac{x^2}{2} + \frac{B}{6}x^3 + \frac{1}{24}x^4$$

$$\begin{aligned}
 l^{(4)}(x) &= A + x + \frac{B}{2}x^2 + \frac{1}{6}x^3 + \frac{C}{24}x^4 \\
 m^{(4)}(x) &= 1 + Bx + \frac{1}{2}x^2 + \frac{C}{6}x^3 + \frac{1}{24}x^4 \\
 n^{(4)}(x) &= B + x + \frac{C}{2}x^2 + \frac{1}{6}x^3 + \frac{D}{24}x^4 \\
 f^{(5)}(x) &= 1 + Ax + \frac{1}{2}x^2 + \frac{B}{6}x^3 + \frac{1}{24}x^4 + \frac{C}{120}x^5 \\
 l^{(5)}(x) &= A + x + \frac{B}{2}x^2 + \frac{1}{6}x^3 + \frac{C}{24}x^4 + \frac{1}{120}x^5 \\
 m^{(5)}(x) &= 1 + Bx + \frac{1}{2}x^2 + \frac{C}{6}x^3 + \frac{1}{24}x^4 + \frac{D}{120}x^5 \\
 f^{(6)}(x) &= 1 + Ax + \frac{1}{2}x^2 + \frac{B}{6}x^3 + \frac{1}{24}x^4 + \frac{C}{120}x^5 + \frac{1}{720}x^6 \\
 l^{(6)}(x) &= A + x + \frac{B}{2}x^2 + \frac{1}{6}x^3 + \frac{C}{24}x^4 + \frac{1}{120}x^5 + \frac{D}{720}x^6 \\
 f^{(7)}(x) &= 1 + Ax + \frac{1}{2}x^2 + \frac{B}{6}x^3 + \frac{1}{24}x^4 + \frac{C}{120}x^5 + \frac{1}{720}x^6 + \frac{D}{5040}x^7 \\
 &\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots
 \end{aligned}$$

Hence, the series solution is given as

$$\begin{aligned}
 f(x) &= 1 + Ax + \frac{1}{2}x^2 + \frac{B}{6}x^3 + \frac{1}{24}x^4 + \frac{C}{120}x^5 + \frac{1}{720}x^6 + \frac{D}{5040}x^7 + \frac{1}{40320}x^8 + \left(\frac{A}{18144} - \frac{1}{362880}\right)x^9 \\
 &\quad + \left(\frac{1}{1209600} - \frac{A}{907200}\right)x^{10} + \left(\frac{A}{6652800} + \frac{B}{1995840} + \frac{1}{5702400}\right)x^{11} \\
 &\quad + \left(\frac{1}{31933440} - \frac{A}{59875200} - \frac{B}{59875200}\right)x^{12} + O(x^{13})
 \end{aligned}$$

Using the boundary conditions at $x = 1$, we get the value of A, B, C and D as

$$A = 0.999870193, B = 1.001257423, C = 0.988438914, D = 1.086357080$$

Finally, the series solution can be written as

$$\begin{aligned}
 f(x) &= 1 + 0.999870193x + \frac{1}{2}x^2 + 0.1668762372x^3 + \frac{1}{24}x^4 + 0.00823699095x^5 + \frac{1}{720}x^6 + 0.000215547x^7 \\
 &\quad + \frac{1}{40320}x^8 + 2.755 \times 10^{-6}x^9 - 2.75 \times 10^{-7}x^{10} + 2.51 \times 10^{-8}x^{11} - 2.1 \times 10^{-9}x^{12} + O(x^{13})
 \end{aligned}$$

Table 1. Comparison of numerical results for example 01.

X	Exact Solution	Approximate Solution	Absolute Error
0	1	1	0
0.1	1.105171	1.105158	0.000013
0.2	1.221403	1.221378	0.000024
0.3	1.349859	1.349825	0.000034
0.4	1.491825	1.491785	0.000039
0.5	1.648721	1.648680	0.000042
0.6	1.822119	1.822079	0.000040
0.7	2.013753	2.013719	0.000034
0.8	2.225541	2.225516	0.000025
0.9	2.459603	2.459590	0.000013
1	2.718282	2.718282	1.66E-10




Figure 1. Comparison of the approximate solution with exact solution for example 01.




Figure 2. Absolute error for example 01.

Example 02 Consider the following 8th order nonlinear boundary value problem

$$\left. \begin{aligned} f^{(8)}(x) &= f(x) - 8xe^x, 0 < x < 1 \\ f(0) &= 1, f^{(2)}(0) = -1, f^{(4)}(0) = -3, f^{(6)}(0) = -5 \\ f(1) &= 0, f^{(2)}(1) = -2e, f^{(4)}(1) = -4e, f^{(6)}(1) = -6e \end{aligned} \right\} \tag{10}$$

The exact solution of this problem is $f(x) = (1 - x)e^x$

The given 8th order boundary value problem can be transformed with the following system

$$\left. \begin{aligned} \frac{df}{dx} &= l(x), \frac{dl}{dx} = m(x), \frac{dm}{dx} = n(x), \frac{dn}{dx} = p(x) \\ \frac{dp}{dx} &= q(x), \frac{dq}{dx} = r(x), \frac{dr}{dx} = s(x), \frac{ds}{dx} = e^{-x}f^2(x) \end{aligned} \right\} \tag{11}$$

With

$$f(0) = 1, l(0) = A, m(0) = -1, n(0) = B, p(0) = -3, q(0) = C, r(0) = -5, s(0) = D$$

The system of differential equation (11) can be written in terms of the following system of integral equations with Lagrange multipliers $\lambda_i = +1; i = 1,2,3, \dots, 8$.

$$\begin{aligned}
 f^{(k+1)}(x) &= 1 + \int_0^x l^{(k)}(t) dt \\
 l^{(k+1)}(x) &= A + \int_0^x m^{(k)}(t) dt \\
 m^{(k+1)}(x) &= -1 + \int_0^x n^{(k)}(t) dt \\
 n^{(k+1)}(x) &= B + \int_0^x p^{(k)}(t) dt \\
 p^{(k+1)}(x) &= -3 + \int_0^x q^{(k)}(t) dt \\
 q^{(k+1)}(x) &= C + \int_0^x r^{(k)}(t) dt \\
 r^{(k+1)}(x) &= -5 + \int_0^x s^{(k)}(t) dt \\
 s^{(k+1)}(x) &= D + \int_0^x (f(t) - 8te^t) dt
 \end{aligned}$$

With

$$f(0) = 1, l(0) = A, m(0) = -1, n(0) = B, p(0) = -3, q(0) = C, r(0) = -5, s(0) = D$$

Now, $f^{(1)}(x) = 1 + Ax$

$$\begin{aligned}
 l^{(1)}(x) &= A - x \\
 m^{(1)}(x) &= -1 + Bx \\
 n^{(1)}(x) &= B - 3x \\
 p^{(1)}(x) &= -3 + Cx \\
 q^{(1)}(x) &= C - 5x \\
 r^{(1)}(x) &= -5 + Dx \\
 s^{(1)}(x) &= D - 8 + x + 8xe^x + 8e^x \\
 f^{(2)}(x) &= 1 + Ax - \frac{x^2}{2} \\
 l^{(2)}(x) &= A - x + \frac{B}{2}x^2 \\
 m^{(2)}(x) &= -1 + Bx - \frac{3}{2}x^2
 \end{aligned}$$

$$\begin{aligned}
 n^{(2)}(x) &= B - 3x + \frac{C}{2}x^2 \\
 p^{(2)}(x) &= -3 + Cx - \frac{5}{2}x^2 \\
 q^{(2)}(x) &= C - x + \frac{D}{2}x^2 \\
 f^{(3)}(x) &= 1 + Ax - \frac{1}{2}x^2 + \frac{B}{6}x^3 \\
 l^{(3)}(x) &= A - x + \frac{B}{2}x^2 - \frac{3}{6}x^3 \\
 m^{(3)}(x) &= -1 + Bx - \frac{3}{2}x^2 + \frac{C}{6}x^3 \\
 n^{(3)}(x) &= B - 3x + \frac{C}{2}x^2 - \frac{5}{6}x^3 \\
 p^{(3)}(x) &= -3 + Cx - \frac{1}{2}x^2 + \frac{D}{6}x^3 \\
 f^{(4)}(x) &= 1 + Ax - \frac{1}{2}x^2 + \frac{B}{6}x^3 - \frac{3}{24}x^4 \\
 l^{(4)}(x) &= A - x + \frac{B}{2}x^2 - \frac{3}{6}x^3 + \frac{C}{24}x^4 \\
 m^{(4)}(x) &= -1 + Bx - \frac{3}{2}x^2 + \frac{C}{3}x^3 - \frac{5}{24}x^4 \\
 n^{(4)}(x) &= B - 3x + \frac{C}{2}x^2 - \frac{1}{6}x^3 + \frac{D}{24}x^4 \\
 f^{(5)}(x) &= 1 + Ax - \frac{1}{2}x^2 + \frac{B}{6}x^3 - \frac{3}{24}x^4 + \frac{C}{120}x^5 \\
 l^{(5)}(x) &= A - x + \frac{B}{2}x^2 - \frac{3}{6}x^3 + \frac{C}{24}x^4 - \frac{5}{120}x^5 \\
 m^{(5)}(x) &= -1 + Bx - \frac{3}{2}x^2 + \frac{C}{3}x^3 - \frac{5}{24}x^4 + \frac{D}{120}x^5 \\
 f^{(6)}(x) &= 1 + Ax - \frac{1}{2}x^2 + \frac{B}{6}x^3 - \frac{3}{24}x^4 + \frac{C}{120}x^5 - \frac{5}{720}x^6 \\
 l^{(6)}(x) &= A - x + \frac{B}{2}x^2 - \frac{3}{6}x^3 + \frac{C}{24}x^4 - \frac{5}{120}x^5 + \frac{D}{720}x^6 \\
 f^{(7)}(x) &= 1 + Ax - \frac{1}{2}x^2 + \frac{B}{6}x^3 - \frac{3}{24}x^4 + \frac{C}{120}x^5 - \frac{5}{720}x^6 + \frac{D}{5040}x^7 \\
 & \vdots
 \end{aligned}$$

Hence, the series solution is given as

$$\begin{aligned}
 f(x) &= 1 + Ax - \frac{1}{2}x^2 + \frac{B}{6}x^3 - \frac{3}{24}x^4 + \frac{C}{120}x^5 - \frac{5}{720}x^6 + \frac{D}{5040}x^7 - \frac{1}{40320}x^8 + \left(\frac{A}{362880} - \frac{1}{45360}\right)x^9 - \frac{1}{403200}x^{10} \\
 & + \left(\frac{B}{39916800} - \frac{1}{4989600}\right)x^{11} - \frac{1}{43545600}x^{12} + O(x^{13})
 \end{aligned}$$

Using the boundary conditions at $x = 1$, we get the value of A, B, C and D as

$$A = 6.772 \times 10^{-7}, B = -2.000006476, C = -3.99994303, D = -6.00036565$$

Finally, the series solution can be written as

$$f(x) = 1 + 6.772 \times 10^{-7}x - 0.50x^2 - 0.3333344127x^3 - \frac{1}{8}x^4 - 0.3333285858x^5 - \frac{1}{144}x^6 - 0.00119054874x^7 - \frac{1}{40320}x^8 - 2.205 \times 10^{-5}x^9 - \frac{1}{403200}x^{10} - 2.505 \times 10^{-7}x^{11} - \frac{1}{43545600}x^{12} + O(x^{13})$$

Table 2. Comparison of numerical results for example 02.

X	Exact Solution	Approximate Solution	Absolute Error
0	1	1	0
0.1	0.994654	0.994676	0.000022
0.2	0.977122	0.977427	0.000305
0.3	0.944901	0.946207	0.001306
0.4	0.895095	0.898480	0.003385
0.5	0.824361	0.830829	0.006468
0.6	0.728848	0.738571	0.009724
0.7	0.604126	0.615376	0.011250
0.8	0.445108	0.452880	0.007772
0.9	0.245960	0.240309	0.005651
1.0	0	-0.035903	0.035903




Figure 3. Comparison of the approximate solution with exact solution for example 02.




Figure 4. Absolute error for example 02.

5. Conclusion

In this paper we have considered eighth order boundary value problem. First, we have shown that fundamentals of variational iteration method. After that we solved two numerical examples by using variational iteration method. then, we show that the approximate solution and compare it with exact solution. We see that, the approximate solution is a series solution and it converse to the exact solution. Finally, we draw a graph of comparison between approximate solution and exact solution and also a graph for absolute error by using Mathematica. Finally, we conclude that the numerical examples show the high degree of efficiency of the variational iteration method.

References

- [1] G. L. Liu, New research directions in the singular perturbation theory: artificial parameter approach and inverse perturbation technique, in: Proceeding of the Conference of the 7th Modern Mathematics and Mechanics, Shanghai, 1997.
- [2] M. A. Noor and S. T. Mohyud-Din, "Homotopy method for solving eighth order boundary value problems," *Journal of Mathematical Analysis and Approximation Theory*, vol. 1, no. 2, pp. 161–169, 2006.
- [3] S. S. Siddiqi and E. H. Twizell, "Spline solution of linear eighth-order boundary value problems," *Computer Methods in Applied Mechanics and Engineering*, vol. 131, no. 3-4, pp. 309–325, 1996.
- [4] A.-M. Wazwaz, "The numerical solution of special eight-order boundary value problems by the modified decomposition method," *Neural, Parallel & Scientific Computations*, vol. 8, no. 2, pp. 133–146, 2000.
- [5] A. Boutayeb and E. H. Twizell, "Finite-difference methods for the solution of special eighth order boundary-value problem," *International Journal of Computer Mathematics*, vol. 48, no. 1, pp. 63–75, 1993.
- [6] R. E. D. Bishop, S. M. Cannon, and S. Miao, "On coupled bending and torsional vibration of uniform beams," *Journal of Sound and Vibration*, vol. 131, no. 1, pp. 457–464, 1989.
- [7] K. Djidjeli, E. H. Twizell, and A. Boutayeb, "Numerical methods for special nonlinear boundary value problems of order $2m$," *Journal of Computational and Applied Mathematics*, vol. 47, no. 1, pp. 35–45, 1993.
- [8] R. P. Agarwal, *Boundary Value Problems for Higher Order Differential Equations*, World Scientific, Teaneck, NJ, USA, 1986.
- [9] A. A. Mamun, M. Asaduzzaman, Solution of seventh order boundary value problem by using variational iteration method, *Int. J. of Mathematics and computational science*, Vol. 5, No. 1, 2019, pp. 6-12, ISSN: 2381-7011 (Print); ISSN: 2381-702X (Online), Published online: April 17, 2019.
- [10] J. H. He, Some asymptotic methods for strongly nonlinear equations, *Int. J. Modern Phys. B* 20 (10) (2006) 1141–1199.
- [11] J. H. He, Non-perturbative methods for strongly nonlinear problems, *Dissertation.de-Verlag im Internet GmbH*, Berlin, 2006.
- [12] J. H. He, Approximate analytical solution for seepage flow with fractional derivatives in porous media, *Comput. Methods Appl. Mech. Engrg.* 167 (1998) 57–68.
- [13] J. H. He, Variational iteration method for autonomous ordinary differential systems, *Appl. Math. Comput.* 114 (2/3) (2000) 115–123.
- [14] J. H. He, X. H. Wu, Construction of solitary solution and compacton-like solution by variational iteration method, *Chaos Solitons Fractals* 29 (1) (2006) 108–113.
- [15] J. H. He, A new approach to nonlinear partial differential equations, *Commun. Nonlinear Sci. Numer. Simul.* 2 (4) (1997) 203–205.
- [16] J. H. He, A variational iteration approach to nonlinear problems and its applications, *Mech. Appl.* 20 (1) (1998) 30–31 (in Chinese).
- [17] J. H. He, Variational iteration method—a kind of nonlinear analytical technique: Some examples, *Int. J. Nonlinear Mech.* 34 (1999) 699–708.
- [18] J. H. He, A generalized variational principle in micromorphic thermoelasticity, *Mech. Res. Commun.* 32 (2005) 93–98.
- [19] J. H. He, Variational iteration method—some recent results and new interpretations, *J. Comput. Appl. Math.* (2006) (in press).
- [20] E. M. Abulwafa, M. A. Abdou, A. A. Mahmoud, The solution of nonlinear coagulation problem with mass loss, *Chaos Solitons Fractals* 29 (2006) 313–330.
- [21] S. Momani, Z. Odibat, Analytical approach to linear fractional partial differential equations arising in fluid mechanics, *Phys. Lett. A* 1 (53) (2006) 1–9.
- [22] S. Momani, S. Abusaad, Application of He's variational-iteration method to Helmholtz equation, *Chaos Solitons Fractals* 27 (5) (2005) 1119–1123.
- [23] M. A. Abdou, A. A. Soliman, Variational iteration method for solving Burgers' and coupled Burgers' equation, *J. Comput. Appl. Math.* 181 (2005) 245–251.
- [24] A. M. Wazwaz, A comparison between the variational iteration method and Adomian decomposition method, *J. Comput. Appl. Math.* (2006) (in press).
- [25] A. M. Wazwaz, The variational iteration method for rational solutions for KdV, K (2,2), Burgers, and cubic Boussinesq equations, *J. Comput. Appl. Math.* (2006) (in press).
- [26] A. M. Wazwaz, *Partial Differential Equations: Methods and Applications*, Balkema Publishers, The Netherlands, 2002.
- [27] A. M. Wazwaz, Necessary conditions for the appearance of noise terms in decomposition solution series, *Appl. Math. Comput.* 81 (1997) 265–274.
- [28] A. M. Wazwaz, A new technique for calculating Adomian polynomials for nonlinear polynomials, *Appl. Math. Comput.* 111 (1) (2000) 33–51.
- [29] A. M. Wazwaz, A new method for solving singular initial value problems in the second order differential equations, *Appl. Math. Comput.* 128 (2002) 47–57.

- [30] A. M. Wazwaz, *A First Course in Integral Equations*, World Scientific, Singapore, 1997.
- [31] A. M. Wazwaz, Analytical approximations and Pad'e approximants for Volterra's population model, *Appl. Math. Comput.* 100 (1999) 13-25.
- [32] G. Adomian, *Solving Frontier Problems of Physics: The Decomposition Method*, Kluwer, Boston, 1994.
- [33] G. Adomian, A review of the decomposition method in applied mathematics, *J. Math. Anal. Appl.* 135 (1988) 501-544.
- [34] J. H. He, Variational principle for some nonlinear partial differential equations with variable coefficients, *Chaos, Solitons & Fractals* 19 (2004) 847-851.
- [35] M. A. Abdou, A. A. Soliman, New applications of variational iteration method, *Physica D* 211 (1-2) (2005) 1-8.
- [36] M. A. Abdou, A. A. Soliman, Variational iteration method for solving Burger's and coupled Burger's equations, *J. comput. Appl. Math.* 181 (2005) 245-251.
- [37] S. Momani, S. Abuasad, Application of He's variational iteration method to Helmholtz equation, *Chaos, Solitons & Fractals* 27 (2006) 1119-1123.
- [38] Z. M. Odibat, S. Momani, Application of variational iteration method to nonlinear differential equations of fractional order, *Int. J. Nonlinear Sci. Numer. Simul.* 7 (2006) 27-36.
- [39] N. Bildik, A. Konuralp, The use of variational iteration method, differential transform method and Adomian decomposition method for solving different types of nonlinear partial differential equations, *Int. J. Nonlinear Sci. Numer. Simul.* 7 (1) (2006) 65-70.
- [40] M. Dehghan, M. Tatar, Identifying an unknown function in a parabolic equation with overspecified data via He's variational iteration method, *Chaos, Solitons and Fractals*, Volume 36, (2008), 157-166.
- [41] M. Dehgha, F. Shakeri, Application of He's variational iteration method for solving the Cauchy reaction-diffusion problem, *Journal of Computational and Applied Mathematics*, Volume 214, (2008), 435-446.
- [42] M. Dehgha, F. Shakeri, Numerical solution of a biological population model using He's variational iteration method, *Computers and Mathematics with Applications*, 54 (2007) 1197-1209.
- [43] M. Dehgha, F. Shakeri, Approximate solution of a differential equation arising in astrophysics using the variational iteration method, *New Astronomy*, 13 (2008) 53-59.